

Contact me in order to access the whole complete document.

Email: solution9159@gmail.com

WhatsApp: <https://wa.me/message/2H3BV2L5TTSUF1>

Telegram: <https://t.me/solutionmanual>

First transform to center-of-mass and relative coordinates.
For $\mu = m/2$, the Schrödinger equation becomes

$$\left[-\frac{\hbar^2}{2\mu} \nabla^2 + \frac{1}{4} K (r-d_0)^2 \right] \psi = E \psi$$

Spherical symmetry allows separation of coordinates, so write $\psi = \frac{1}{r} R(r) Y_{\ell m}(\theta, \phi)$. The radial component is then

$$\left[-\frac{\hbar^2}{2\mu} \frac{d^2}{dr^2} + \frac{\hbar^2 \ell(\ell+1)}{2\mu r^2} + \frac{1}{4} K (r-d_0)^2 \right] R(r) = E R(r)$$

Put $K \equiv 2\mu\omega_0^2$; $\frac{\hbar^2 \ell(\ell+1)}{2\mu d_0^4} \equiv \gamma_2 K$; $r-d_0 \equiv \rho$

Expand to second order in ρ and drop terms in γ_2^2 since $\gamma_2 \ll 1$

$$\left\{ -\frac{\hbar^2}{2\mu} \frac{d^2}{dr^2} + \frac{1}{2} \mu \omega_0^2 \left[(1+12\gamma_2)(\rho-4\gamma_2 d_0)^2 + 4\gamma_2 d_0^2 \right] \right\} R = E R$$

The energy levels of this simple harmonic oscillator are

$$E_{\ell n} = \left(n + \frac{1}{2}\right) \hbar \omega_0 (1+6\gamma_2) + \gamma_2 K d_0^2 + O(\gamma_2^2)$$

So $E_{00} = \frac{1}{2} \hbar \sqrt{\frac{K}{m}}$ $E_{01} = \frac{3}{2} \hbar \sqrt{\frac{K}{m}}$

$$E_{10} = \frac{1}{2} \hbar \sqrt{\frac{K}{m}} (1+6\gamma_1) + K d_0^2 \gamma_1$$

$$E_{11} = \frac{3}{2} \hbar \sqrt{\frac{K}{m}} (1+6\gamma_1) + K d_0^2 \gamma_1$$

Then

$$(E_{11} - E_{00}) - (E_{01} - E_{00}) - (E_{10} - E_{00})$$

$$= 6 \hbar \sqrt{\frac{K}{m}} \gamma_1 = \boxed{\frac{12 \hbar^3}{d_0^4 \sqrt{K m^3}}}$$

Solution to 1.2

Incoming wave is $E_0 e^{i(kx - \omega_0 t)}$ with $\omega_0 = kc$.

Atom is at $x = \sum_q a_q \sin(\omega_q t + \phi_q)$

$$\begin{aligned} S_0 E &= E_0 e^{ik \sum_q a_q \sin(\omega_q t + \phi_q) - i\omega_0 t} \\ &= E_0 e^{-i\omega_0 t} \left[1 + \frac{i\omega_0}{c} \sum_q a_q \sin(\omega_q t + \phi_q) \right. \\ &\quad \left. - \frac{\omega_0^2}{2c^2} \sum_{q, q'} a_q a_{q'} \sin(\omega_q t + \phi_q) \sin(\omega_{q'} t + \phi_{q'}) \right] \end{aligned}$$

with higher terms neglected if $kx \ll 1$. In the sum over q and q' , the phases ϕ_q are random, and so these terms average to zero unless $q = q'$, in which case $\sum a_q a_{q'} \sin(\omega_q t + \phi_q) \sin(\omega_{q'} t + \phi_{q'})$ becomes

$$\sum_q a_q^2 \langle \sin^2(\omega_q t + \phi_q) \rangle = \frac{1}{2} \sum_q a_q^2.$$

$$\begin{aligned} \text{Thus } E(t) &= e^{-i\omega_0 t} \left(1 - \frac{\omega_0^2}{4c^2} \sum_q a_q^2 \right) \\ &\quad \pm \sum_q e^{-i(\omega_0 \pm \omega_q)t \pm i\phi_q} \frac{\omega_0}{2c} a_q \end{aligned}$$


Intensity is square of amplitude.

We have an unshifted peak of height

$$1 - 2 \frac{\omega_0^2}{4c^2} \sum_q a_q^2 \approx \exp\left(-\frac{\omega_0^2}{2c^2} \sum_q a_q^2\right)$$

and side bands of intensity $\propto \frac{\omega_0^2}{4c^2} |a_q|^2$.

Note: $\exp\left(-\frac{\omega_0^2}{4c^2} \sum_q a_q^2\right)$ is known as the Debye-Waller factor.



Solution to 1.3

(i) Assume longitudinal wave, \vec{q} in z direction

$$\sum_{\ell} \vec{E}_{\ell} = \sum_{\ell} E_{\ell} \hat{z} = \sum_{\ell} z e y_0 e^{iq\ell \cos\theta} (3\cos^2\theta - 1) \ell^{-3/2}$$

For small q replace sum by integral

$$\sum_{\ell} E_{\ell} \rightarrow \frac{z e y_0}{\Omega} \iint e^{iq\ell \cos\theta} (3\cos^2\theta - 1) \frac{2\pi}{\ell} d\ell \sin\theta d\theta$$

vol. of unit cell

Put $I \equiv \int e^{iq\ell \cos\theta} \sin\theta d\theta = \frac{2 \sin q\ell}{q\ell} = \frac{2 \sin x}{x}$, say.

$$\begin{aligned} \text{Then } \Sigma E &= - \frac{2\pi z e y_0}{\Omega} \int_{q\ell}^{qR} (I + 3 \frac{d^2 I}{dx^2}) \frac{dx}{x} \\ &= - \frac{2\pi z e y_0}{\Omega} \int \left[\frac{2 \sin x}{x} + 3 \left(-\frac{2 \sin x}{x} - \frac{4 \cos x}{x^2} + \frac{4 \sin x}{x^3} \right) \right] \frac{dx}{x} \\ &= - \frac{2\pi z e y_0}{\Omega} \int 4 \left[-\frac{\sin x}{x^2} - \frac{3 \cos x}{x^3} + \frac{3 \sin x}{x^4} \right] dx \\ &= \frac{8\pi z e y_0}{\Omega} \left[\frac{\sin x - x \cos x}{x^3} \right]_{q\ell}^{qR} \end{aligned}$$

Alternative way: say $e^{iq \cdot \vec{r}} \sim \sum (2\ell+1) i^{\ell} j_{\ell}(qr) P_{\ell}(\cos\theta)$
 and $3\cos^2\theta - 1 \propto P_2(\cos\theta)$.
 Thus $\Sigma E \propto \int x^{-3} j_2(qr) x^2 dx \propto \left[\frac{j_1(qr)}{qr} \right]$

So if $q\ell \rightarrow 0$ and $qR \rightarrow \infty$

$$\begin{aligned} \Sigma E &= \frac{8\pi z e y_0}{\Omega} \left(\frac{x - x^{3/6} + \dots - x + x^{3/2} - \dots}{x^3} \right)_{0}^{\infty} \\ &= \frac{8\pi z e y_0}{\Omega} \times \frac{1}{3} \end{aligned}$$

If $M \ddot{y}_{\ell} = z e \Sigma E_{\ell}$, then $\omega^2 = \frac{2}{3} \times \frac{4\pi (ze)^2}{M \Omega} = \boxed{\frac{2}{3} \omega_p^2}$


soln. to 1.3 (cont.)

but if $qR \rightarrow 0$, then $\omega^2 = 0$

(ii) Cylinder

$$\Sigma E = \frac{2\pi z e y_0}{\Omega} \int_0^R e^{iqz} \left[\frac{3z^2}{(\rho^2+z^2)^{5/2}} - \frac{1}{(\rho^2+z^2)^{3/2}} \right] \rho d\rho dz$$

$$= \frac{2\pi z e y_0}{\Omega} \int_0^R e^{iqz} dz \left[\frac{\rho^2}{(\rho^2+z^2)^{3/2}} \right]_0^R$$



For lower limit $qE \rightarrow 0$, and integral is easy. For upper limit, put $z = R \tan \theta$

$$dz = R \sec^2 \theta d\theta$$

$$\int e^{iqz} dz \frac{R^2}{(R^2+z^2)^{3/2}} = \int e^{iqR \tan \theta} \frac{R^3 \sec^3 \theta d\theta}{R^3 \sec^3 \theta}$$

$$= \int e^{iqR \tan \theta} \cos \theta d\theta$$

Now the question is, what are the limits of θ ?

If $z_{\max} \gg R$, then limits of θ are $\pm \pi/2$, and then integral is 2 if $qR \rightarrow 0$ and 0 if $qR \rightarrow \infty$.

If $R \gg z_{\max}$, limits of θ are small, and integral vanishes.

So what's going on? Explanation lies in depolarizing factors.

(i) Sphere, $qR \rightarrow 0$

$E = 0$ inside hole

Sphere $qR \rightarrow \infty$

Outer surface cancels $E \neq 0$ at center as inner surface acts.

(ii) Cylinder $z_{\max} \gg R$

$qR \rightarrow 0$
 $E = 0$

$R \gg z_{\max}$

$E \neq 0$

Problem 1.4. Phonon interactions

Assume a linear dispersion relation, so that $\omega = v|\vec{q}|$.

Then incoming wave has amplitude $\propto \exp(i\vec{q}_1 \cdot \vec{r} - i\omega_1 t)$

and outgoing wave $\propto \exp(i\vec{q}_3 \cdot \vec{r} - i\omega_3 t)$.

The diffraction grating is moving with velocity \vec{v} , so transform to coordinates moving with the grating.

Then put $\vec{r}' \equiv \vec{r} - \vec{v}t$

$$\phi_{\text{incoming}} \propto \exp i[\vec{q}_1 \cdot (\vec{r}' + \vec{v}t) - \omega_1 t]$$

$$\phi_{\text{outgoing}} \propto \exp i[\vec{q}_3 \cdot (\vec{r}' + \vec{v}t) - \omega_3 t]$$

At points spaced a distance \vec{l} along the grating, the phases must be matched, so

$$i[\vec{q}_1 \cdot (\vec{l} + \vec{v}t) - \omega_1 t] = i[\vec{q}_3 \cdot (\vec{l} + \vec{v}t) - \omega_3 t] + 2\pi n \quad \text{integer } n$$

So (A) $\vec{q}_1 \cdot \vec{v} - \omega_1 = \vec{q}_3 \cdot \vec{v} - \omega_3$

and (B) $\vec{q}_1 \cdot \vec{l} = \vec{q}_3 \cdot \vec{l} - 2\pi$ for first-order diffraction

Now $\vec{v} = \frac{\omega_2 \vec{q}_2}{q_2^2}$ and $\vec{l} = \frac{2\pi \vec{q}_2}{q_2^2}$, so from (B)

$$\vec{q}_1 \cdot \vec{q}_2 = \vec{q}_3 \cdot \vec{q}_2 - \vec{q}_2 \cdot \vec{q}_2 \quad \text{and by symmetry also}$$


$$\vec{q}_2 \cdot \vec{q}_1 = \vec{q}_3 \cdot \vec{q}_1 - \vec{q}_1 \cdot \vec{q}_1$$

Try $\vec{q}_3 = \vec{q}_1 + \vec{q}_2 + \vec{k}$. Then $\vec{k} \cdot \vec{q}_2 = 0$ and $\vec{k} \cdot \vec{q}_1 = 0$ so $\vec{k} = 0$.

From (A) $\frac{\vec{q}_1 \cdot \vec{q}_2}{q_2^2} \omega_2 - \omega_1 = \frac{(\vec{q}_1 + \vec{q}_2) \cdot \vec{q}_2}{q_2^2} \omega_2 - \omega_3$

and so $\boxed{\omega_3 = \omega_1 + \omega_2}$ when $\boxed{\vec{q}_3 = \vec{q}_1 + \vec{q}_2}$

This is the classical analog of energy and momentum conservation for phonons.



Problem 1.5 Anharmonic chain

Equations of motion are $M\ddot{y}_n = 4g[(y_{n+1} - y_n)^3 - (y_n - y_{n-1})^3]$

Put $r_n = y_n - y_{n-1}$. Then $M\ddot{r}_n = 4g[r_{n+1}^3 - 2r_n^3 + r_{n-1}^3]$

Call lattice spacing = a . For a traveling wave $r_n = A f(na - vt)$.

Here A is an amplitude and f is dimensionless

$$\text{Now } \ddot{r}_n = \frac{Av^2}{a^2} \frac{\partial^2 f}{\partial n^2} \text{ and } \frac{MAv^2}{a^2} \frac{\partial^2 f}{\partial n^2} = 4gA^3 (f_{n+1}^3 - 2f_n^3 + f_{n-1}^3)$$

$$\text{Put } h \equiv \sqrt{\frac{4gA^2a^2}{Mv^2}} f. \text{ Then } \frac{\partial^2 h}{\partial n^2} = h_{n+1}^3 - 2h_n^3 + h_{n-1}^3$$

This equation has no parameters. Suppose it has a solution \tilde{h}_n . Then $r_n = \sqrt{\frac{Mv^2}{4ga^2}} \tilde{h}$, and

$$(r_n)_{\max} = \frac{v}{2a} \sqrt{\frac{M}{g}} (\tilde{h})_{\max} \quad \text{so } \boxed{v \propto (r_n)_{\max}}$$

A simpler solution comes from dimensional analysis
Velocity v must be proportional to the lattice spacing,
so try $v = \text{constant} \times a \times g^\alpha M^\beta (r_{\max})^\gamma$

$$\left[\frac{\text{length}}{\text{time}} \right] = [\text{length}] \times \left[\frac{\text{mass}}{\text{length}^2 \text{time}^2} \right]^\alpha \times [\text{mass}]^\beta [\text{length}]^\gamma$$

$$\alpha = \frac{1}{2}; \quad \beta = -\frac{1}{2}; \quad \gamma = 1 \quad \text{so } \boxed{v \propto r_{\max}}$$

Problem 1.6 Suppose original Fermi radius k_F is reduced to k_{\downarrow} for down-spin electrons and increased to k_{\uparrow} for up-spin electrons. By conservation of the number of electrons $k_{\uparrow}^3 + k_{\downarrow}^3 = 2k_F^3$

so we can put $k_{\uparrow}^3 = (1+\alpha)k_F^3$ and $k_{\downarrow}^3 = (1-\alpha)k_F^3$

Magnetic field shifts the Fermi energies to make

$$\frac{\hbar^2}{2m} (k_{\uparrow}^2 - k_{\downarrow}^2) = 2\mu_B H \text{ so } E_F [(1+\alpha)^{2/3} - (1-\alpha)^{2/3}] = 2\mu_B H$$

For small α , $(1 + \frac{2}{3}\alpha - \frac{1}{9}\alpha^2 \dots) - (1 - \frac{2}{3}\alpha - \frac{1}{9}\alpha^2 \dots) = \frac{2\mu_B H}{E_F}$

$$\text{so } \alpha \approx \frac{3\mu_B H}{2E_F}$$

We were given the change in total kinetic energy
Now total kinetic energy is proportional to k_F^5 , as energy $\propto k_F^2$ and number of electrons $\propto k_F^3$.

$$\text{Thus } \frac{k_{\uparrow}^5 + k_{\downarrow}^5}{2k_F^5} = 1 + 5 \times 10^{-8}$$

$$\frac{1}{2} [(1+\alpha)^{5/3} + (1-\alpha)^{5/3}] \approx 1 + \frac{5/3 \times 2/3}{2} \alpha^2 = 1 + \frac{5}{9} \alpha^2$$

$$\text{So } \alpha^2 = 9 \times 10^{-8} \text{ and } \alpha = 3 \times 10^{-4}$$

Hence
$$H = 2 \times 10^{-4} \frac{E_F}{\mu_B}$$

1.7 Solution $E_{kinetic} \propto N_{\uparrow} k_{\uparrow}^2 + N_{\downarrow} k_{\downarrow}^2 \propto k_{\uparrow}^5 + k_{\downarrow}^5$
 $E_{int} \propto N_{\uparrow}^{4/3} + N_{\downarrow}^{4/3} \propto k_{\uparrow}^4 + k_{\downarrow}^4$.


Thus $E_{total} = a(k_{\uparrow}^5 + k_{\downarrow}^5) + b(k_{\uparrow}^4 + k_{\downarrow}^4)$

Minimize this subject to the constraint $k_{\uparrow}^3 + k_{\downarrow}^3 = 2k_F^3$ (A)

Lagrange undetermined multiplier λ for constraint gives
 $5a k_{\uparrow}^4 + 4b k_{\uparrow}^3 - 3\lambda k_{\uparrow}^2 = 0$; $5a k_{\downarrow}^4 + 4b k_{\downarrow}^3 - 3\lambda k_{\downarrow}^2 = 0$

Eliminate λ to find $5a(k_{\uparrow}^2 - k_{\downarrow}^2) + 4b(k_{\uparrow} - k_{\downarrow}) = 0$

Hence $k_{\downarrow} = k_{\uparrow}$ or $k_{\uparrow} + k_{\downarrow} = -4b/5a$.



Plot a graph of the constraint (A).

Question is: Where on this line is

E_{tot} a minimum? If b is positive,

then by inspection of the expression for E_{tot} we see that $k_{\uparrow} = k_{\downarrow}$ is a minimum, so magnetism is unstable. This remains true until line $k_{\uparrow} + k_{\downarrow} = -4b/5a$ intersects the constraint curve. Then \rightarrow

But if there is a maximum on this

curve, then magnetism can be

metastable, since variation of E_{tot} with k_{\uparrow} can show that there is an energy barrier separating magnetized from unmagnetized state. This intersection occurs when

$k_{\downarrow} = 0$ $k_{\uparrow} = 2^{1/3} k_F = -4b/5a$. As b becomes

more negative, we reach the point where

$E_{tot}(k_{\uparrow} = k_{\downarrow}) = E_{tot}(k_{\uparrow} = 0)$. This occurs

when $2ak_F^5 + 2bk_F^4 = a2^{5/3}k_F^5 + b2^{4/3}k_F^4$, from which

$b = -(2^{1/3} + 1)a k_F$ Beyond this point the magnetized state is stable.




Now $E_{kin}(k_{\uparrow} = k_{\downarrow}) = \frac{3}{5} N \epsilon_F$, so $a = \frac{3}{5} N \epsilon_F / 2k_F^5$.

$E_{int}(k_{\uparrow} = k_{\downarrow}) = 2K(N/2)^{4/3}$, so $b = K(N/2)^{4/3} / k_F^4$

So stable if $K < -(2^{1/3} + 1) 2^{1/3} 3\epsilon_F / 5N^{1/3}$ i.e. $K < -1.708 \epsilon_F / N^{1/3}$

unstable if $K > -3\epsilon_F / 2^{4/3} N^{1/3}$ i.e. $K > -1.191 \epsilon_F / N^{1/3}$

metastable in-between.



Problem 1.8 Antiferromagnetic magnons.

Analysis similar to that of section 1.4 leads to substitutions
 $\vec{\mu}^{(a)} = \vec{\mu}_0 + \vec{\mu}_1 e^{i(\omega t + \vec{k} \cdot \vec{r})}$; $\vec{\mu}_1 = \vec{\mu}_0 + \vec{\mu}_1 e^{i(\omega t + \vec{k} \cdot \vec{r})}$

Linearized equations of motion are

$$i\omega \vec{\mu}_1^{(a)} = C \sum_{\text{neighbors}} (\vec{\mu}_0^{(b)} \times \vec{\mu}_1^{(a)} + \vec{\mu}_1^{(b)} \times \vec{\mu}_0^{(a)} e^{i\vec{k} \cdot (\vec{r}' - \vec{r})})$$

$$i\omega \vec{\mu}_1^{(b)} = C \sum (\vec{\mu}_0^{(a)} \times \vec{\mu}_1^{(b)} + \vec{\mu}_1^{(a)} \times \vec{\mu}_0^{(b)} e^{i\vec{k} \cdot (\vec{r}' - \vec{r})})$$

Now put $\sum e^{i\vec{k} \cdot (\vec{r}' - \vec{r})} \equiv \sum f(\vec{k})$ with \sum = coordination number

$$\begin{aligned} \text{Then } (C\sum\mu_0 - \omega)\mu_1^{(a)} + C\sum\mu_0 f(\vec{k})\mu_1^{(b)} &= 0 \\ -C\sum\mu_0 f(\vec{k})\mu_1^{(a)} - (C\sum\mu_0 + \omega)\mu_1^{(b)} &= 0 \end{aligned}$$

$$\text{From which } \omega^2 = (C\sum\mu_0)^2 (1 - f^2(\vec{k}))$$

$$\text{Put } C\sum\mu_0 \equiv \omega_{\text{max}} \quad \boxed{\omega = \omega_{\text{max}} \sqrt{1 - f^2(\vec{k})}}$$

For simple cubic lattice $f(\vec{k}) = \frac{1}{3} [\cos k_x a + \cos k_y a + \cos k_z a]$

In (111) direction $\vec{k} = k(1, 1, 1)$ and $f(k) = \cos ka$

$$\sqrt{1 - f^2} = \sin ka$$



$$\frac{\mu_1^{(a)}}{\mu_1^{(b)}} = \frac{-f}{1 \pm \sqrt{1 - f^2}} = -\frac{\omega_{\text{max}} + \omega}{\omega_{\text{max}} - \omega} \quad \text{or} \quad -\frac{\omega_{\text{max}} - \omega}{\omega_{\text{max}} + \omega}$$

so for small ω , $\mu_1^{(a)} \approx -\mu_1^{(b)}$

for large ω

$$\frac{\mu_1^{(a)}}{\mu_1^{(b)}} \rightarrow 0 \text{ or } \infty$$



Problem 1.9

$$\Delta E_{\text{magnetic}} = -\mu_B H (N_{\uparrow} - N_{\downarrow}) = -\mu_B H \frac{N}{2} \left(\frac{k_{\uparrow}^3 - k_{\downarrow}^3}{k_F^3} \right)$$

With $k_{\uparrow}^3 = (1+\alpha)k_F^3$, $\Delta E_{\text{magnetic}} = -N\mu_B H \alpha$

$$E_{\text{kinetic}} = \frac{3}{5} N \epsilon_F \left(\frac{k_{\uparrow}^5 + k_{\downarrow}^5}{2k_F^5} \right) \approx \frac{3}{5} N \epsilon_F \left(1 + \frac{5}{9} \alpha^2 \right)$$

Also $\epsilon_F \left[(1+\alpha)^{2/3} - (1-\alpha)^{2/3} \right] = 2\mu_B H$, so $\alpha \approx \frac{3\mu_B H}{2\epsilon_F}$.

Thus $\Delta E_{\text{total}} = \Delta E_{\text{kinetic}} + \Delta E_{\text{magnetic}}$

$$= \frac{3}{5} N \epsilon_F \left[\frac{5}{9} \left(\frac{3\mu_B H}{2\epsilon_F} \right)^2 - \frac{3(\mu_B H)^2}{\frac{3}{5}\epsilon_F \times 2\epsilon_F} \right] = \frac{3}{5} N \epsilon_F \left[-\frac{5}{4} \left(\frac{\mu_B H}{\epsilon_F} \right)^2 \right]$$

So $-\frac{5}{4} \left(\frac{\mu_B H}{\epsilon_F} \right)^2 = -5 \times 10^{-8}$ and $H = 2 \times 10^{-4} \frac{\epsilon_F}{\mu_B}$ as before.

Quick solution:

Total energy is of the form $ax^2 - bx$ where $x \propto N_{\uparrow} - N_{\downarrow}$.

Minimum of this expression is at $x = \frac{b}{2a}$

$$\text{Then } \Delta(\text{kinetic energy}) = ax^2 = \frac{b^2}{4a}$$

$$\Delta(\text{potential energy}) = -bx = -\frac{b^2}{2a}$$

So $\Delta(\text{total energy}) = -\Delta(\text{kinetic energy})$

and thus H must be the same as in problem 1.6

Problem 1.10 Energy of a Toda soliton.

From Eq. (1.3.2) $z_n = -\frac{1}{b} \ln [1 + \sinh^2 \mu \operatorname{sech}^2(\mu n - \beta t)]$

Now $1 + \sinh^2 \mu \operatorname{sech}^2(\mu n - \beta t) = \frac{\cosh^2(\mu n - \beta t) + \sinh^2 \mu}{\cosh^2(\mu n - \beta t)}$

$$= \frac{\cosh^2(\mu n - \beta t) (\cosh^2 \mu - \sinh^2 \mu) + \sinh^2 \mu (\cosh^2(\mu n - \beta t) - \sinh^2(\mu n - \beta t))}{\cosh^2(\mu n - \beta t)}$$

$$= \frac{\cosh[\mu(n+1) - \beta t] \cosh[\mu(n-1) - \beta t]}{\cosh^2(\mu n - \beta t)}$$

Define $A_n \equiv \frac{\cosh(\mu n - \beta t)}{\cosh(\mu(n-1) - \beta t)}$ so $z_n = -\frac{1}{b} \ln \left(\frac{A_{n+1}}{A_n} \right)$

or $z_n = \frac{1}{b} (\ln A_n - \ln A_{n+1})$ so $\sum_{-\infty}^{\infty} z_n = \frac{1}{b} (\ln A_0 - \ln A_{-\infty}) = -2\mu/b$

The energy of the soliton is

$$\mathcal{E} = \sum_{-\infty}^{\infty} \left[\frac{m}{2} \dot{y}_n^2 + \frac{a}{b} (e^{-\beta y_n} - 1) + a z_n \right]$$

Now \dot{y}_n can be found from the equation $y_n - y_{-\infty} = \sum_{-\infty}^n z_n$

and $\sum_{-\infty}^n z_n = -\frac{1}{b} (\ln A_{n+1} - \ln A_{-\infty})$

so $\dot{y}_n = -\frac{1}{b} \frac{d}{dt} (\ln A_{n+1}) = \frac{\beta}{b} [\tanh(\mu(n+1) - \beta t) - \tanh(\mu n - \beta t)]$

The energy is constant, so we can put $t=0$, and $\beta = \sqrt{\frac{ab}{m}} \sinh \mu$

$$\mathcal{E} = \sum_{-\infty}^{\infty} \frac{a}{2b} \sinh^2 \mu \left\{ (\tanh \mu(n+1) - \tanh \mu n)^2 + 2 \operatorname{sech}^2 \mu n \right\} - \frac{2\mu a}{b}$$

$$= \frac{a}{2b} \sinh^2 \mu \sum_{-\infty}^{\infty} \left[\tanh^2 \mu(n+1) - \tanh^2 \mu n + 2(1 - \tanh \mu n \tanh \mu(n+1)) \right]$$

$$= \frac{a}{b} \sinh^2 \mu \sum_{-\infty}^{\infty} (1 - \tanh \mu n \tanh \mu(n+1)) - \frac{2\mu a}{b}$$

Prob. 1.10 continued

$$\text{Now } \tanh \mu = \tanh[(n+1)\mu - n\mu] = \frac{\tanh(n+1)\mu - \tanh n\mu}{1 - \tanh(n+1)\mu \tanh n\mu}$$

$$\text{so } 1 - \tanh n\mu \tanh \mu(n+1) = \coth \mu (\tanh(n+1)\mu - \tanh n\mu)$$

$$\begin{aligned} \text{and } E &= \frac{a}{b} \sinh^2 \mu \coth \mu \sum (\tanh(n+1)\mu - \tanh n\mu) - \frac{2\mu a}{b} \\ &= \frac{a}{b} \sinh \mu \coth \mu \times 2 - \frac{2\mu a}{b} \end{aligned}$$

$$E = \frac{2a}{b} (\sinh \mu \coth \mu - \mu)$$

(Thanks to Xin-Yi Wang for this formulation)

Easy version: For large μ at $t=0$, $r_0 \sim -\frac{1}{b} \ln(\sinh^2 \mu)$
so $r_0 \sim -\frac{2\mu}{b}$ and $r_{\pm 1} \sim -\frac{1}{b} \ln 2 \ll r_0$

Potential energy $V \sim ar_0 + \frac{a}{b} e^{-br_0} \sim \frac{a}{b} \sinh^2 \mu$.

Also $\dot{r}_0 = 0$ $\dot{r}_{\pm 1} \sim \frac{\beta}{b} (\pm 1) \sim \pm \sqrt{\frac{a}{bM}} \sinh \mu$

and kinetic energy $\sim \frac{a}{b} \sinh^2 \mu$ so $E_{\text{large } \mu} \sim \frac{2a}{b} \sinh^2 \mu$

For small μ at $t=0$ the motion is harmonic, and so potential energy = kinetic energy. We just need to calculate the kinetic energy and double it.

Now $\dot{r}_n \sim \frac{\beta}{\mu} \frac{\partial r_n}{\partial n}$ and thus $\dot{y}_n \sim \frac{\beta}{\mu} r_n$ so $\frac{1}{2} m \dot{y}_n^2 \sim \frac{m \beta^2 r_n^2}{2\mu^2}$

$$\begin{aligned} \sum \frac{1}{2} m \dot{y}_n^2 &\sim \frac{a}{2b} \frac{\sinh^2 \mu}{\mu^2} \sum_n [\sinh^2 \mu \operatorname{sech}^2 \mu n]^2 \\ &\sim \frac{a\mu^3}{2b} \int_{-\infty}^{\infty} \operatorname{sech}^4 x dx \sim \frac{2a\mu^3}{3b} \text{ so } E_{\text{small } \mu} \sim \frac{a\mu^3}{b} \end{aligned}$$